

Probing mSUGRA Models at Linear Colliders^a

T. Kamon,^b R. Arnowitt, B. Dutta,^c and V. Khotilovich

Department of Physics, Texas A&M University, College Station, TX 77843-4242, USA

Abstract

A feasibility study of a 500-GeV linear collider is presented for mSUGRA models in co-annihilation region. We find an active mask is critical to suppress $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ events to probe the models.

1 Feasibility of 500-GeV Linear Collider

The grand unification of the three gauge coupling constants arise naturally in minimal SUGRA (mSUGRA) models with universal soft breaking parameters ($m_{1/2}$, m_0 , A_0 , $\tan\beta$, sign of μ).^{1,2} The experimental data to provide the most significant constraints are the Higgs mass bound, the $b \rightarrow s\gamma$ branching ratio, and (possibly) the muon magnetic moment anomaly (a_μ) along with the amount of relic density of the lightest neutralino ($\tilde{\chi}_1^0$) as cold dark matter. These constraints produce a lower bound of $m_{1/2} \gtrsim 300$ GeV (Higgs mass and $b \rightarrow s\gamma$) and an upper bound of $m_{1/2} \lesssim 900$ GeV (a_μ), making the SUSY spectrum well accessible to the LHC.³ The relic density constraints limit the most part of the allowed parameter space to narrow band where $\tilde{\tau}_1\tilde{\chi}_1^0$ co-annihilation occurs.^{4,5} See, for example, Fig. 1.⁶ Further, at large $\tan\beta$ (e.g., 40), $M_{\tilde{\tau}_1}$ becomes the next to lightest SUSY particle and all other SUSY particles would be too heavy to produce in pair at e^+e^- linear colliders (LCs). The only accessible production at 500-GeV LCs could be $\tilde{\chi}_2^0\tilde{\chi}_1^0$, $\tilde{\tau}_1^+\tilde{\tau}_1^-$ and $\tilde{\chi}_1^0\tilde{\chi}_1^0$, where $M_{\tilde{\chi}_1^\pm} > 250$ GeV and $\Delta M \equiv M_{\tilde{\tau}_1} - M_{\tilde{\chi}_1^0} = 5 - 30$ GeV. Furthermore, the $\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1\tau$ decay is dominant. This guides us to the conclusion that one of the most critical experimental signatures is $\tau^+\tau^-$ plus missing energy. We present a feasibility study of a 500-GeV LC for the signature of two hadronically-decaying τ lepton (τ_h) plus missing energy.

^aT.K. and V.K. are supported by DOE grant DE-FG03-95ER40917, and R.A. and B.D. by NSF grant PHY-0101015.

^be-mail address: t-kamon@tamu.edu

^cPresent Address: *Department of Physics, University of Regina, Regina SK, S4S 0A2, Canada*




Figure 1: Parameter space, in m_0 - $m_{1/2}$ plane of mSUGRA models ($\tan\beta = 40$), allowed by relic density of $\tilde{\chi}_1^0$'s as cold dark matter.⁶ Also shown are branching ratios for $B_s \rightarrow \mu^+\mu^-$, where three dashed lines indicate $Br = 2 \times 10^{-8}$, 1×10^{-8} , and 7×10^{-9} . The Tevatron Run II with 15 fb^{-1} will be accessible to 1×10^{-8} . The vertical dotted lines label Higgs masses. The three short solid lines indicate the $\sigma_{p\tilde{\chi}_1^0}$ values (from left: $0.03 \times 10^{-6} \text{ pb}$, $0.002 \times 10^{-6} \text{ pb}$, $0.001 \times 10^{-6} \text{ pb}$).

We generate (i) $\tilde{\chi}_2^0\tilde{\chi}_1^0$ and $\tilde{\tau}_1^+\tilde{\tau}_1^-$ events via ISAJET and (ii) SM four-fermion final states of $\nu_{e,\mu,\tau}\bar{\nu}_{e,\mu,\tau}\tau^+\tau^-$ from weak process and $e^+e^-\tau^+\tau^-$ ($p_{T\tau} > 3 \text{ GeV}/c$) from two-photon ($\gamma\gamma$) process via WPHACT.⁷ For the SUSY events, we choose three mSUGRA points, as listed in Table 1. Those events are simulated and analyzed with the LCDRoot software package.⁸ The electron-beam polarization is set to be $\mathcal{P}(e^-) = 0.9$ (L.H) to enhance the $\tilde{\chi}_2^0\tilde{\chi}_1^0$ signal and the opposite polarization $\mathcal{P}(e^-) = -0.9$ (R.H) is used for the $\tilde{\tau}_1^+\tilde{\tau}_1^-$ signal extraction. See Table 2.

For the L.H. case, we optimize the selection cuts for $\tilde{\chi}_2^0\tilde{\chi}_1^0$ events over the $\tilde{\tau}_1^+\tilde{\tau}_1^-$ and SM events: (1) $N_{jet} = 2$ (JADE Y-cut ≥ 0.0025), where each jet passes a τ_h identification (1 or 3 tracks; $M_{jet} < 2 \text{ GeV}/c^2$) with $E_{jet} > 3 \text{ GeV}$; (2) Opposite charge between two jets ($q_{j1} \cdot q_{j2} = -1$); (3) $-q_j \cdot \cos\theta_j < 0.75$; (4) Missing p_T ($\equiv \not{p}_T$) $> 5, 10$, or $20 \text{ GeV}/c$; (5) $\cos\theta(\vec{p}_{j2}, \vec{p}_{vis}) > -0.6$; (6) Acoplanarity $> 40^\circ$; (7) Vetoing event if there is any EM cluster with $E_{cluster} > 2 \text{ GeV}$ in $0.995 < |\cos\theta_{cluster}| < 0.9$ ($5.8^\circ < \theta_{cluster} < 25.8^\circ$), or any cluster with $E_{cluster} > 100 \text{ GeV}$.

Table 1: Three mSUGRA points for $\mu > 0$, $\tan \beta = 40$, and $A_0 = 0$ by ISAJET.⁷

	$m_{1/2}$	m_0	$M_{\tilde{\chi}_2^0}$	$M_{\tilde{\tau}_1}$	$M_{\tilde{\chi}_1^0}$	ΔM ($\equiv M_{\tilde{\tau}_1} - M_{\tilde{\chi}_1^0}$)
A1	360	210	266	149.9	144.2	5.7
A2	360	215	266	154.8	144.2	10.6
A3	360	225	266	164.3	144.4	20.2

Table 2: Production cross sections (fb) for the SM (excluding $\gamma\gamma$ process) and SUSY events in e^+e^- collisions with different electron-beam polarizations ($\mathcal{P}(e^-) = -0.9, 0$, and 0.9) at $\sqrt{s} = 500$ GeV.

$\mathcal{P}(e^-)$	-0.9 (R.H.)		0		+0.9 (L.H.)		
SM (excluding $\gamma\gamma$)	7.84		48.9		89.8		
SUSY		$\tilde{\chi}_1^0\tilde{\chi}_2^0$	$\tilde{\tau}_1\tilde{\tau}_1$	$\tilde{\chi}_1^0\tilde{\chi}_2^0$	$\tilde{\tau}_1\tilde{\tau}_1$	$\tilde{\chi}_1^0\tilde{\chi}_2^0$	$\tilde{\tau}_1\tilde{\tau}_1$
	A1	0.529	26.4	3.39	19.6	7.10	12.8
	A2	0.520	24.4	3.31	18.4	6.91	11.8
	A3	0.501	21.1	3.15	15.8	6.62	10.3

in $2(1)^\circ < \theta_{cluster} < 5.8^\circ$, or electron/positron with $E_e > 1.5$ GeV in $|\cos \theta_e| < 0.9$. In the present study, the same cuts are used for the R.H. case. The results are summarized in Table 3. With an active mask of covering down to 2° , the LC detector would be able to test mSUGRA models with small ΔM , and being better with 1° mask.

2 Conclusion

We have examined a feasibility of a 500-GeV LC in probing three mSUGRA points in co-annihilation region. Our preliminary result showed that an active mask is very critical for the study, to suppress $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ events. In future study, we will include $e^+e^- \rightarrow e^+e^-q\bar{q}$ where the quark jets are misidentified as τ_h 's.

References

1. A. H. Chamseddine, R. Arnowitt, and P. Nath, Phys. Rev. Lett. **49**, 970 (1982).

Table 3: Number of events expected at 500-GeV LC with 500 fb^{-1} . Three mSUGRA points are examined. Boldfaced numbers for L.H. (R.H.) electron-beams indicate $> 5\sigma$ -discovery sensitivities for $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ ($\tilde{\tau}_1^+ \tilde{\tau}_1^-$) over $\tilde{\tau}_1^+ \tilde{\tau}_1^- + \text{SM}$ ($\tilde{\chi}_2^0 \tilde{\chi}_1^0 + \text{SM}$) background events. The SM “weak” process includes all $\nu_{e,\mu,\tau} \bar{\nu}_{e,\mu,\tau} \tau^+ \tau^-$ final states, while the SM $\gamma\gamma$ is the $ee\tau\tau$ ($p_{T\tau} > 3 \text{ GeV}/c$) final state.

Process		$\mathcal{P}(e^-) = 0.9(\text{L.H.})$			$\mathcal{P}(e^-) = -0.9(\text{R.H.})$		
		$p_T^{\min} = 5$	10	20	5	10	20
$\tilde{\chi}_2^0 \tilde{\chi}_1^0$	A1	609	545	402	48	42	31
	A2	952	873	626	71	65	47
	A3	1180	1113	835	90	85	63
$\tilde{\tau}_1^+ \tilde{\tau}_1^-$	A1	170	16	0	353	37	0
	A2	687	394	30	1442	817	59
	A3	1198	994	438	2510	2098	939
SM weak		2486	2320	1797	391	373	312
SM $\gamma\gamma$	2-5.8° mask	1507	13	0	1507	13	0
	[1-5.8° mask]	[38]	[2]	[0]	[38]	[2]	[0]
	[no mask]	[11400]	[3020]	[89]	[11400]	[3020]	[89]

2. R. Barbieri, S. Ferrara, and C. A. Savoy, Phys. Lett. B**119**, 343 (1982); L. J. Hall, J. Lykken, and S. Weinberg, Phys. Rev. D**27**, 2359 (1983); P. Nath, R. Arnowitt, and A. H. Chamseddine, Nucl. Phys. B**227**, 121 (1983).
3. See, for example, R. Arnowitt and B. Dutta, arXiv:hep-ph/0211042.
4. K. Griest and D. Seckel, Phys. Rev. D**43**, 3191 (1991); P. Gondolo and G. Gelmini, Nucl. Phys. B**360**, 145 (1991).
5. J. Ellis, T. Falk, K. Olive, M. Srednicki, Astropart. Phys. **13**, 181 (2000); Erratum-ibid. **15**, 413 (2001); J. Ellis, T. Falk, G. Ganis, K. Olive, M. Srednicki, Phys. Lett. B**510**, 236 (2001); R. Arnowitt, B. Dutta, and Y. Santos, hep-ph/0010244; Nucl. Phys. B**606**, 59 (2001); M. Gomez and J. Vergados, hep-ph/0012020; M. Gomez, G. Lazarides, C. Pallis, Phys. Rev. D**61**, 123512 (2000); Phys. Lett. B**487**, 313 (2000).
6. R. Arnowitt, B. Dutta, T. Kamon, and M. Tanaka, Phys. Lett. B**538**, 121 (2002).
7. We use ISAJET v7.63 (H. Baer *et al.*), Tauola v2.6, WPHACT v2.02pol (A. Ballestrero *et al.*; all four-fermion final states with e^+/e^- polarization).
8. LCDRoot Package v3.5 with LD Mar01 detector parametrization of EM and HAD calorimeters ($|\cos\theta| < 0.995$) and tracking system ($|\cos\theta| < 0.9$).